

Exclusive $B \rightarrow M\nu\bar{\nu}$ ($M = \pi, K, \rho, K^*$) Decays and Leptophobic Z' Model

Jong Hun Jeon^{1*}, C. S. Kim^{1†}, Jake Lee^{2‡} and Chaehyun Yu^{3§}

¹ *Department of Physics, Yonsei University, Seoul 120-479, Korea*

² *Department of Physics, Sogang University, Seoul 100-611, Korea*

³ *Department of Physics, Korea University, Seoul 136-701, Korea*

(Dated: February 2, 2008)

Abstract

We consider the exclusive flavor changing neutral current processes $B \rightarrow M\nu\bar{\nu}$ ($M = \pi, K, \rho, K^*$) in the leptophobic Z' model, in which the charged leptons do not couple to the extra Z' boson. We find that these exclusive modes are very effective to constrain the leptophobic Z' model. In the leptophobic Z' model, additional right-handed neutrinos are introduced and they can contribute to the missing energy signal in $B \rightarrow M + \cancel{E}$ decays. Through the explicit calculations, we obtain quite stringent bounds on the model parameters, $|U_{sb}^{Z'}| \leq 0.29$ and $|U_{db}^{Z'}| \leq 0.61$, from the already existing experimental data. We also briefly discuss an interesting subject of massive right-handed neutrinos, which might be connected with the dark matter problem.

* jhjeon@cskim.yonsei.ac.kr

† cskim@yonsei.ac.kr

‡ jilee@cskim.yonsei.ac.kr

§ chyu@korea.ac.kr

I. INTRODUCTION

Flavor changing neutral current (FCNC) processes have gained continuous interests as powerful probes of new physics beyond the standard model (SM) as well as stringent tests within the SM. Since FCNC processes are only generated at the loop level within the SM, they are suppressed by the loop momentum and off-diagonal Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. Therefore, if FCNC exists in the tree level or exists with an enhancement factor in the coupling without severe suppression, it would be very sensitive to new physics effects beyond the SM. B -factory experiments such as Belle and BaBar have given various opportunities to test those FCNC processes. Among the various terms in the effective Hamiltonian responsible for the decay of b quark, the FCNCs are represented by QCD penguin and electroweak (EW) penguin operators. There are plenty of QCD or EW penguin dominant processes in non-leptonic and semi-leptonic B decays. As an example, recent experimental data for QCD penguin dominant $B \rightarrow K\pi$ decays appear to be very interesting: Branching ratios and direct and mixing-induced CP asymmetries could not be consistently explained within the current SM frameworks, for example, QCD factorization (QCDF) [1], perturbative QCD (PQCD) [2], and soft-collinear effective theory (SCET) models [3]. Although it has been claimed in Ref. [4] that all the current data except for the mixing-induced CP asymmetry could be accommodated within the PQCD scheme by considering the next-leading-order corrections, there is still room for new physics or new mechanism beyond the SM, especially, in the EW penguin sector [5, 6, 7] to explain all the experimental data simultaneously.

The EW penguin contributions have been thoroughly studied in the SM and beyond, too: While the EW penguin effect with the photon emission in $b \rightarrow s\gamma$ processes has been precisely determined, and already put stringent bounds on the relevant new physics scenarios, experimental measurements for the Z -mediated EW penguin effect are now under way at B factories through $B \rightarrow X\ell\ell$ and $B \rightarrow X\nu\bar{\nu}$ decay processes. Although it is experimentally very difficult to measure, $B \rightarrow X\nu\bar{\nu}$ decay is an extremely good channel for the study of Z -mediated EW penguin contribution within the SM and beyond [8, 9, 10, 11, 12].

In many new physics scenarios, an extra $U(1)$ gauge boson is usually introduced after the high energy symmetry breaking or as a low energy effective theory [13, 14, 15]. In

general, the extra gauge boson, commonly named as Z' boson, has a possibility to induce the FCNCs. In the present work, we focus on the E_6 -based leptophobic Z' model [16] among various possibilities with the extra $U(1)$ gauge boson extension such as the string-inspired grand unified theories (GUTs) [13, 17], dynamical symmetry breaking models [18], string inspired models [19], extra dimension models [20], and little higgs models [21]. It is well known that the extra neutral gauge boson can be leptophobic by introducing the kinetic mixing term in the E_6 -based model. Originally, the leptophobic Z' model was introduced to explain the R_b - R_c puzzle at LEP [22] and anomalous high- E_T jet cross section at CDF [23]. Although its original motivation has been disappeared, the leptophobic Z' model certainly remains as a viable scenario beyond the SM. The main experimental constraints on the (non-leptophobic) Z' boson arise from the electroweak precision measurements at LEP or direct searches via Drell-Yan processes at Tevatron. For the leptophobic Z' boson, however, we have now only one experimental constraint from the dijet experiments at D0 which exclude the mass range for $365 \text{ GeV} \leq M_{Z'} \leq 615 \text{ GeV}$ [24]. For the future experiments, while the CERN Large Hadron Collider (LHC) could serve as a probe of the leptophobic Z' boson via $pp \rightarrow \gamma + \cancel{E}$, the future e^-e^+ linear collider (ILC) may have no role as far as leptophobia is concerned. In this regard, we note that $B \rightarrow M + \cancel{E}$ decays, where M is a pseudoscalar or a vector boson, will be a good channel for the probe of the leptophobic Z' model.

In this paper, we investigate $B \rightarrow M\nu\bar{\nu}$ ($M = \pi, K, \rho, K^*$) decays in the leptophobic Z' model as a possible candidate of new physics in the EW penguin sector. First, we briefly introduce the leptophobic Z' model based on the E_6 grand unified theory (GUT) in Sec. II. Sec. III deals with the effective Hamiltonian and decay rates in detail. In Sec. IV we present numerical analysis and discuss its implications. Concluding remarks are also in Sec. IV.

II. LEPTOPHOBIC Z' MODEL AND FCNC

First, we briefly review the leptophobic Z' model. In GUT or string-inspired point of view, the E_6 model is a very plausible extension of the SM [25]. It is natural that a $U(1)'$ gauge group remains as a low energy effective theory after the symmetry breaking of the E_6 group. We assume that the E_6 group is broken through the following breaking chain

$$\begin{aligned} E_6 &\rightarrow SO(10) \times U(1)_\psi \\ &\rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi \end{aligned}$$

$$\rightarrow SU(2)_L \times U(1) \times U(1)', \quad (1)$$

where $U(1)'$ is a linear combination of two additional $U(1)$ gauge groups with

$$Q' = Q_\psi \cos \theta - Q_\chi \sin \theta,$$

where θ is the familiar E_6 mixing angle.

The most general Lagrangian, which is invariant under the SM gauge group with an extra $U(1)'$, allows the kinetic mixing term $\mathcal{L}_{\text{mixing}} = -\frac{\sin \chi}{2} \tilde{B}_{\mu\nu} \tilde{Z}'^{\mu\nu}$ between the $U(1)$ and $U(1)'$ gauge boson fields. This off-diagonal term can be removed by the non-unitarity transformation

$$\tilde{B}_\mu = B_\mu - \tan \chi Z'_\mu, \quad \tilde{Z}'_\mu = \frac{Z'_\mu}{\cos \chi}, \quad (2)$$

which leads to the possibility of leptophobia of the physical Z' gauge boson with the E_6 mixing. Once all the couplings are GUT normalized, the interaction Lagrangian of fermion fields and Z' gauge boson can be written as

$$\mathcal{L}_{\text{int}} = -\lambda \frac{g_2}{\cos \theta_W} \sqrt{\frac{5 \sin^2 \theta_W}{3}} \bar{\psi} \gamma^\mu \left(Q' + \sqrt{\frac{3}{5}} \delta Y_{SM} \right) \psi Z'_\mu, \quad (3)$$

where the ratio of gauge couplings $\lambda = g_{Q'}/g_Y$, and $\delta = -\tan \chi/\lambda$ [25]. The general fermion- Z' couplings depend on two free parameters, $\tan \theta$ and δ , effectively [26]. The Z' boson can be leptophobic when $(Q' + \sqrt{\frac{3}{5}} \delta Y_{SM}) = 0$ for L and e^c simultaneously. L and e^c are the lepton fields in the **27** representation of E_6 , as shown in Table I. In the conventional embedding, the Z' boson can be made leptophobic with $\delta = -1/3$ and $\tan \theta = \sqrt{3/5}$. In other embedded schemes, where the quantum numbers between L and H (or e^c , ν^c , and S^c) are switched, one can find suitable values for δ and $\tan \theta$ which make the Z' boson leptophobic [16].

In the leptophobic Z' model, FCNCs can arise through the mixing between the ordinary SM fermions and exotic fermions. Since the mixing of the left-handed fermions may lead to the relatively large Z -mediated FCNCs, we only allow the mixing of the right-handed fermions. That is, the mixing between $d_R(s_R, b_R)$ and h_R can induce the FCNCs when their Z' charges are different [16]. After integrating out degrees of freedom of heavy exotic fermions and gauge bosons, the FCNC Lagrangian for the $b \rightarrow q(q = s, d)$ transition can be parameterized as

$$\mathcal{L}_{\text{FCNC}}^{Z'} = -\frac{g_2}{2 \cos \theta_W} U_{qb}^{Z'} \bar{q}_R \gamma^\mu b_R Z'_\mu, \quad (4)$$

TABLE I: Charges of fermions contained in the **27** representation of E_6 within the conventional embedding [25].

Particle	$SU(3)_c$	Y	$2\sqrt{6}Q_\psi$	$2\sqrt{10}Q_\chi$
$Q = (u, d)^T$	3	1/6	1	-1
$L = (\nu, e)^T$	1	-1/2	1	3
u^c	$\bar{3}$	-2/3	1	-1
d^c	$\bar{3}$	1/3	1	3
e^c	1	1	1	-1
ν^c	1	0	1	-5
$H = (N, E)^T$	1	-1/2	-2	-2
$H^c = (N^c, E^c)^T$	1	1/2	-2	2
h	3	-1/3	-2	2
h^c	$\bar{3}$	1/3	-2	-2
S^c	1	0	4	0

where all the theoretical uncertainties including the mixing parameters are absorbed into the coupling $U_{qb}^{Z'}$. Then, the experimental constraints on $U_{qb}^{Z'}$ may be obtained by considering related non-leptonic B decays such as $B \rightarrow K\pi$ decays for which one may also find a clue to a solution of the *recently-popular* $B \rightarrow K\pi$ puzzle. But, it is not easy to obtain reliable bounds on $U_{qb}^{Z'}$ because the non-leptonic B decays severely depend on unknown strong phases [27] and, furthermore, the EW penguin amplitude is sub-dominant in most non-leptonic decays.

It is obvious that if new physics effects are mainly through the EW penguin contributions, the EW penguin dominant process can give the most powerful constraints, and it is natural to consider the processes such as $b \rightarrow s\gamma, b \rightarrow s(d)\ell\ell, b \rightarrow s(d)\nu\bar{\nu}$ as appropriate probes of the relevant new physics. In the most new physics scenarios such as the minimal supersymmetric standard model (MSSM), model parameters are connected to many different observables, and therefore, in order to get consistent constraints on the model parameters we have to consider all the relevant processes simultaneously. As we will see later, however, one can consider only $b \rightarrow s(d)\nu\bar{\nu}$ decays for the leptophobic Z' model and almost free from other experimental constraints.

III. THEORETICAL DETAILS FOR $B \rightarrow M\nu\bar{\nu}$ ($M = \pi, K, \rho, K^*$) DECAYS

In the SM, the effective Hamiltonian describing $b \rightarrow q\nu\bar{\nu}$ ($q = d, s$) decays is given by

$$H_{\text{eff}}(b \rightarrow q\nu_{\text{SM}}\bar{\nu}_{\text{SM}}) = \frac{G_F\alpha}{2\pi\sqrt{2}}V_{tb}V_{tq}^*C_{10}^\nu\bar{q}\gamma^\mu(1-\gamma^5)b\bar{\nu}\gamma_\mu(1-\gamma^5)\nu, \quad (5)$$

where G_F is the Fermi constant, α is the fine structure constant and V_{ij} are elements of the CKM matrix. The Wilson coefficient C_{10}^ν is dominated by the short-distance dynamics associated with top quark exchange [9], and has the theoretical uncertainty due to the error of top quark mass, whose explicit form can be found in literatures [10, 28, 29]. Therefore, aside from the CKM matrix elements, for the $b \rightarrow q\nu\bar{\nu}$ decays, there remains only well-controllable theoretical uncertainty from C_{10}^ν . Although it is not easy to measure experimentally due to only single meson with the missing energy in the final state, it can be analyzed through the reconstruction of another B meson from the $\Upsilon(4S) \rightarrow B^+B^-$ event.

Because the leptophobia means that the $U(1)'$ charge is zero for all the ordinary left-handed and right-handed lepton fields within the SM, the extra leptophobic Z' gives no additional contribution to $B \rightarrow M + \cancel{E}$ decays if the missing energy is solely due to the SM neutrinos. However, ν^c or S^c in the **27** representation of E_6 , as shown in Table I, may be interpreted as the right-handed neutrino and so there can arise additional contributions to the missing energy signal in $B \rightarrow M + \cancel{E}$ decays.

Here we take ν^c as a candidate for the right-handed neutrino and assume that other heavy exotic fermions and bosons be integrated out. Then, the effective Hamiltonian responsible for $b \rightarrow q\nu_R\bar{\nu}_R$ decay is written as

$$H_{\text{eff}}(b \rightarrow q\nu_R\bar{\nu}_R) = \frac{\pi\alpha}{\sin^2 2\theta_W M_{Z'}^2}U_{qb}^{Z'}Q_{\nu_R}^{Z'}\bar{q}\gamma^\mu(1+\gamma_5)b\bar{\nu}\gamma_\mu(1+\gamma_5)\nu, \quad (6)$$

where

$$Q_{\nu_R}^{Z'} = \frac{1}{2}\kappa\sqrt{\frac{5\sin^2\theta_W}{3}}, \quad (7)$$

and κ depends on the details of unification. Here we take $\kappa = 1$ for simplicity. Then, the total branching ratios (BRs) for $B \rightarrow M + \cancel{E}$ signals are given by

$$\mathcal{B}(B \rightarrow M\nu\bar{\nu}) = \mathcal{B}(B \rightarrow M\nu_{\text{SM}}\bar{\nu}_{\text{SM}}) + \mathcal{B}(B \rightarrow M\nu_R\bar{\nu}_R). \quad (8)$$

First we take all the neutrinos massless, then the differential decay rates for $B \rightarrow M\nu\bar{\nu}$ ($M = P, V$) in the leptophobic Z' model are given by

$$\frac{d\Gamma}{dq^2}(B^\pm \rightarrow P^\pm\nu\bar{\nu}) = \frac{G_F^2\alpha^2 M_B^3}{2^8\pi^5}|V_{tb}V_{tq}^*|^2|C_{10}^\nu|^2$$

$$\times \left[1_{\text{SM}} + \frac{\pi^2 Q_{\nu_R}^{Z'}{}^2 |U_{qb}^{Z'}|^2}{\alpha^2 |V_{tb} V_{tq}^*|^2 |C_{10}^\nu|^2} \left(\frac{M_Z^2}{M_{Z'}^2} \right)^2 \right] \lambda_P^{3/2} |f_+^P(q^2)|^2, \quad (9)$$

$$\begin{aligned} \frac{d\Gamma}{dq^2}(B^\pm \rightarrow V^\pm \nu \bar{\nu}) &= \frac{G_F^2 \alpha^2 M_B^3}{2^{10} \pi^5} |V_{tb} V_{tq}^*|^2 |C_{10}^\nu|^2 \lambda_V^{1/2} \\ &\times \left[1_{\text{SM}} + \frac{\pi^2 Q_{\nu_R}^{Z'}{}^2 |U_{qb}^{Z'}|^2}{\alpha^2 |V_{tb} V_{tq}^*|^2 |C_{10}^\nu|^2} \left(\frac{M_Z^2}{M_{Z'}^2} \right)^2 \right] \\ &\times \left[\frac{8 \lambda_V \tilde{q}^2}{(1 + \sqrt{r_V})^2} V^V(q^2)^2 + \frac{1}{r_V} \left\{ (1 + \sqrt{r_V})^2 (\lambda_V + 12 r_V \tilde{q}^2) A_1^V(q^2)^2 \right. \right. \\ &\left. \left. + \frac{\lambda_V^2}{(1 + \sqrt{r_V})^2} A_2^V(q^2)^2 - 2 \lambda_V (1 - r_V - \tilde{q}^2) \text{Re} (A_1^V(q^2) A_2^V(q^2)^*) \right\} \right] \quad (10) \end{aligned}$$

where $\lambda_M = 1 + r_M^2 + \tilde{q}^4 - 2r_M - 2\tilde{q}^2 - 2r_M \tilde{q}^2$ with $r_M = M_M^2/M_B^2$, $\tilde{q}^2 = q^2/M_B^2$ and $P(V)$ in the superscript and subscript stands for a pseudoscalar (vector) meson, respectively.

Main theoretical uncertainties arise from the hadronic transition form factors for the B meson decay into the $P(V)$ meson. However, one can also reduce these uncertainties by taking ratios of the BRs, related by $SU(3)$ flavor symmetry or isospin symmetry, such as $B \rightarrow \pi(\rho) \nu \bar{\nu}$ and $B \rightarrow \pi(\rho) \ell \nu$ [10]. Especially, the corrections to the strict isospin symmetry due to mass difference between $\rho^\pm$ and ρ^0 and form factors are expected to be very small.

IV. NUMERICAL ANALYSIS AND DISCUSSIONS

In Table II, we show the theoretical estimates of BRs within the SM and their current experimental bounds at B factories [30, 31]. The errors of the SM estimates in Tab. II are mainly due to the hadronic transition form factors and the CKM matrix elements: For the form factors, we use recent results from the LCSR model with theoretical uncertainties of about 10 to 13 % at zero momentum transfer [32, 33]:

$$f_+^\pi(0) = 0.258 \pm 0.031, \quad f_+^K(0) = 0.331 \pm 0.041, \quad (11)$$

$$V^\rho(0) = 0.323 \pm 0.030, \quad V^{K^*}(0) = 0.411 \pm 0.033, \quad (12)$$

$$A_1^\rho(0) = 0.242 \pm 0.023, \quad A_1^{K^*}(0) = 0.292 \pm 0.028, \quad (13)$$

$$A_2^\rho(0) = 0.221 \pm 0.023, \quad A_2^{K^*}(0) = 0.259 \pm 0.027. \quad (14)$$

TABLE II: Expected BRs in the SM and experimental bounds (90% C.L.) in units of 10^{-6} .

mode	BRs in the SM	Experimental bounds
$B \rightarrow K\nu\bar{\nu}$	$5.31^{+1.11}_{-1.03}$	< 36 [30]
$B \rightarrow \pi\nu\bar{\nu}$	$0.22^{+0.27}_{-0.17}$	< 100 [31]
$B \rightarrow K^*\nu\bar{\nu}$	$11.15^{+3.05}_{-2.70}$	-
$B \rightarrow \rho\nu\bar{\nu}$	$0.49^{+0.61}_{-0.38}$	-

The details of the form factors are given in Appendix. For the CKM matrix elements, we adopted PDG results [34],

$$|V_{tb}| \simeq 1, \quad |V_{ts}| = 0.040 \pm 0.003, \quad |V_{td}| = (9.4 \pm 4.6) \times 10^{-3}. \quad (15)$$

Please note that $b \rightarrow d$ transition modes receive much larger theoretical uncertainties from $|V_{td}|$, as can be seen in Table II. The BRs of $B \rightarrow K^{(*)}\nu\bar{\nu}$ decays, which are the $b \rightarrow s$ transition process, are of an order of 10^{-5} to 10^{-6} . On the other hand, the BRs of $B \rightarrow \pi(\rho)\nu\bar{\nu}$ arising from the $b \rightarrow d$ transition are less by about one order. And as expected from larger degrees of freedom in final states, BRs of the vector boson production processes are about 2 or 3 times larger than the corresponding pseudoscalar processes.

Recently the Belle and BaBar Collaborations presented the upper limits on BRs of $B \rightarrow K\nu\bar{\nu}$ and $B \rightarrow \pi\nu\bar{\nu}$ decays [30, 31]. For the K meson production process with the missing energy the current experimental bound is about 7 times larger than the SM expectation, whereas for the π meson production the upper bound is much higher by an order of 10^3 than the theoretical estimate. Therefore, there is plenty of room for new physics effects in these decays at present. The BRs of all the modes are expected to be (more precisely) measured at B factories soon.

Now we investigate additional effects from the leptophobic Z' boson. In Figs. 1 and 2, we present our predictions for the BRs in the leptophobic Z' model as a function of the effective coupling $|U_{qb}^{Z'}|$ in the model. The solid and dotted lines denote the SM predictions and the current experimental bounds, respectively. In the leptophobic Z' model, we have two parameters, the mass of Z' boson and new FCNC coupling, $|U_{qb}^{Z'}|$. Since the D0 experiment excludes the mass range $365 \text{ GeV} \leq M_{Z'} \leq 615 \text{ GeV}$ [24], we take $M_{Z'} = 700 \text{ GeV}$, which is also consistent with the mass bound of the conventional non-leptophobic Z' model. Although we choose a specific mass for the Z' boson, our analysis can be easily translated through the




FIG. 1: Branching ratios for (a) $B^\pm \rightarrow K^\pm \nu \bar{\nu}$ and (b) $B^\pm \rightarrow \pi^\pm \nu \bar{\nu}$, where ν can be the ordinary SM neutrinos or right-handed neutrinos.




FIG. 2: Branching ratios for (a) $B^\pm \rightarrow K^{*\pm} \nu \bar{\nu}$ and (b) $B^\pm \rightarrow \rho^\pm \nu \bar{\nu}$, where ν can be the ordinary SM neutrinos or right-handed neutrinos.

corresponding changes in the effective coupling $|U_{qb}^{Z'}|$ for different Z' masses. Considering the current experimental upper bounds, one can extract the following constraints for the FCNC couplings from Fig. 1,

$$|U_{sb}^{Z'}| \leq 0.29, \quad |U_{db}^{Z'}| \leq 0.61, \quad (16)$$

for $B \rightarrow K \nu \bar{\nu}$ and $B \rightarrow \pi \nu \bar{\nu}$ decays, respectively. Compared with the inclusive $b \rightarrow s \nu \bar{\nu}$ decay case [16], we find presently the exclusive mode gives more stringent bounds on the

leptophobic FCNC couplings.

As already mentioned, although the exclusive $B \rightarrow M\nu\bar{\nu}$ decay modes are much easier at the experimental detection than the inclusive ones, they have inevitable theoretical uncertainties from hadronic form factors. For example, our expectation value for $B \rightarrow K\nu\bar{\nu}$ in the SM with recent form factors from LCSR model is about 5.31×10^{-6} , however, it can be changed to $\sim 3.8 \times 10^{-6}$ by using different form factors [9]. In order to reduce the form factor uncertainties, we take ratios for $\mathcal{B}(B \rightarrow M\nu\bar{\nu})$ to $\mathcal{B}(B \rightarrow Me\nu)$ for $M = \pi, \rho$ mesons. In the limit of isospin symmetry, and requiring $M_\pi^\pm = M_\pi^0, M_\rho^\pm = M_\rho^0$, we obtain

$$\begin{aligned} \frac{\mathcal{B}(B^\pm \rightarrow \pi^\pm \nu\bar{\nu})}{\mathcal{B}(B^\pm \rightarrow \pi^0 e^\pm \nu)} &= \frac{3\alpha^2}{2\pi^2} \left| \frac{V_{td}}{V_{ub}} \right|^2 |C_{10}^\nu|^2 + \frac{3}{2} \frac{Q_{\nu_R}^{Z'^2} |U_{db}^{Z'}|^2}{|V_{ub}|^2} \left(\frac{M_Z^2}{M_{Z'}^2} \right)^2 \\ &= 2.5 \times 10^{-3} \left(\frac{0.40}{|V_{ub}/V_{td}|} \right)^2 + 7.6 \times 10^{-3} \left(\frac{3.7 \times 10^{-3}}{|V_{ub}|} \right)^2 \left(\frac{700 \text{ GeV}}{M_{Z'}} \right)^4 \left(\frac{|U_{db}^{Z'}|}{0.05} \right)^2, \quad (17) \end{aligned}$$

$$\begin{aligned} \frac{\mathcal{B}(B^\pm \rightarrow \rho^\pm \nu\bar{\nu})}{\mathcal{B}(B^\pm \rightarrow \rho^0 e^\pm \nu)} &= \frac{3\alpha^2}{2\pi^2} \left| \frac{V_{td}}{V_{ub}} \right|^2 |C_{10}^\nu|^2 + \frac{3}{2} \frac{Q_{\nu_R}^{Z'^2} |U_{db}^{Z'}|^2}{|V_{ub}|^2} \left(\frac{M_Z^2}{M_{Z'}^2} \right)^2 \\ &= 2.5 \times 10^{-3} \left(\frac{0.40}{|V_{ub}/V_{td}|} \right)^2 + 7.6 \times 10^{-3} \left(\frac{3.7 \times 10^{-3}}{|V_{ub}|} \right)^2 \left(\frac{700 \text{ GeV}}{M_{Z'}} \right)^4 \left(\frac{|U_{db}^{Z'}|}{0.05} \right)^2, \quad (18) \end{aligned}$$

where the form factor dependencies are all cancelled out. For $K^{(*)}$ meson productions, however, since there does not exist such a corresponding semi-leptonic decay mode, one cannot adopt the same analysis given above. Instead, considering $M_\rho \approx M_{K^*}$ and the flavor $SU(3)$ symmetry, we obtain the following relation:

$$\begin{aligned} \frac{\mathcal{B}(B^\pm \rightarrow K^{*\pm} \nu\bar{\nu})}{\mathcal{B}(B^\pm \rightarrow \rho^0 e^\pm \nu)} &\simeq \frac{3\alpha^2}{2\pi^2} \left| \frac{V_{ts}}{V_{ub}} \right|^2 |C_{10}^\nu|^2 + \frac{3}{2} \frac{Q_{\nu_R}^{Z'^2} |U_{sb}^{Z'}|^2}{|V_{ub}|^2} \left(\frac{M_Z^2}{M_{Z'}^2} \right)^2 \\ &= 4.5 \times 10^{-2} \left(\frac{9.3 \times 10^{-2}}{|V_{ub}/V_{ts}|} \right)^2 + 1.2 \times 10^{-1} \left(\frac{3.7 \times 10^{-3}}{|V_{ub}|} \right)^2 \left(\frac{700 \text{ GeV}}{M_{Z'}} \right)^4 \left(\frac{|U_{sb}^{Z'}|}{0.2} \right)^2. \quad (19) \end{aligned}$$

On the contrary to $B^\pm \rightarrow K^* \nu\bar{\nu}$ and $B^\pm \rightarrow \rho^0 e^\pm \nu$, we cannot use the $SU(3)$ flavor symmetry between $B^\pm \rightarrow K^\pm \nu\bar{\nu}$ and $B^\pm \rightarrow \pi^0 e^\pm \nu$ transitions because of the possibly large $SU(3)$ breaking due to large mass difference between π and K mesons.

Up to now we have assumed all the neutrinos are massless. Aside from the SM neutrinos of which masses are indeed very small, the extra right-handed neutrinos in the leptophobic Z' model can have sizable masses. Effects of the massive neutrinos will be an interesting subject by themselves. Here we briefly introduce the mass effects. In Fig. 3, we present the differential BRs as a function of the normalized momentum transfer-squared for $m_{\nu_R} =$




FIG. 3: Differential BRs as a function of the normalized momentum transfer square, $\tilde{q}^2 = q^2/M_B^2$, in units of 10^{-6} , for (a) $B^\pm \rightarrow K^\pm \nu \bar{\nu}$, (b) $B^\pm \rightarrow \pi^\pm \nu \bar{\nu}$, (c) $B^\pm \rightarrow K^{*\pm} \nu \bar{\nu}$, and (d) $B^\pm \rightarrow \rho^\pm \nu \bar{\nu}$. Here the decay rates in the leptophobic Z' model are normalized to be five times larger than those in the SM.

0, 0.2, 0.5, 1 GeV. We normalized the integrated BRs such that the BRs in the leptophobic Z' model are five times larger than those in the SM. It is apparent that the differential BRs show the specific features according to the corresponding mass of the right-handed neutrino. Especially the sharp rise near the threshold point allows to increase the accuracy of the mass measurement of the lightest right-handed neutrino. Unfortunately, if its mass is lower than a few hundred MeV, it is very hard to find the difference from the massless neutrino case.

In general, the additional right-handed neutrinos can have Dirac mass terms coupled with

the ordinary SM neutrinos as well as Majorana masses [35]. Since there is no limit for both Dirac and Majorana masses in principle, they are free parameters. One interesting scenario is a so-called “ ν MSM” model [36], which allows three right-handed (light) neutrinos in addition to the SM ones. Assuming that Majorana masses of right-handed neutrinos are the order of the electroweak scale or below and Yukawa couplings are very small, this model can be accommodated with the present neutrino data. Especially, if masses of the right-handed neutrinos are in range of $2 \lesssim M_{\nu_R} \lesssim 5$ keV, they can be good candidates for the warm dark matters, and they can explain the neutrino oscillation via the see-saw mechanism and baryon asymmetry of the universe as well [36].

To conclude, the FCNC processes with the missing energy signal in B decays have been studied in the leptophobic Z' model. In general the direct probe of the leptophobic Z' model may be very difficult at ILC as well as LHC because of large hadronic backgrounds. In this regard the exclusive FCNC processes $B \rightarrow M\nu\bar{\nu}$ are very adequate to measure new physics in the leptophobic Z' model with the right-handed neutrinos. This model could be quite important in the context of possibly large new physics scenario in the EW penguin sector. We also showed that ratios of BRs can reduce the large hadronic uncertainty from form factors. The differential BRs are very useful if the right-handed neutrinos have the sub-GeV masses. We also noted that the right-handed neutrinos could be accommodated with the “ ν MSM” scenario.

ACKNOWLEDGMENTS

We thank Y.G. Kim and U. Haisch for careful reading and useful comments. The work of J.H.J. was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) No. R02-2003-000-10050-0. The work of C.S.K. was supported in part by CHEP-SRC Program and in part by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) No. KRF-2005-070-C00030. The work of J.L. was supported in part by Brain Korea 21 Program and in part by Grant No. F01-2004-000-10292-0 of KOSEF-NSFC International Collaborative Research Grant. The work of C.Y. was supported by the Korea Research Foundation Grant funded by Korea Government (MOEHRD, Basic Research Promotion FUND) (KRF-2005-075-C00008).

TABLE III: The parameters for form factors in LCSR model [32, 33] where $m_b = 4.8$ GeV is fixed.

parameter	r_1	r_2	m_{fit}^2	m_R^2
f_+^K	0.1616	0.1730	—	5.41^2
f_0^K	—	0.3302	37.46	—
f_+^π	0.744	-0.486	40.73	5.32^2
f_0^π	—	0.258	33.81	—
V^{K^*}	0.923	-0.511	49.40	5.32^2
$A_0^{K^*}$	1.364	-0.990	36.78	5.28^2
$A_1^{K^*}$	—	0.290	40.38	—
$A_2^{K^*}$	-0.084	0.342	52.00	—
V^ρ	1.045	-0.721	38.34	5.32^2
A_0^ρ	1.527	-1.220	33.36	5.28^2
A_1^ρ	—	0.240	37.51	—
A_2^ρ	0.009	0.212	40.82	—

APPENDIX : HADRONIC FORM FACTORS

The hadronic matrix elements for $B \rightarrow P\nu\bar{\nu}$ ($P = \pi, K$) decays can be parameterized in terms of the form factors $f_+^P(q^2)$ and $f_-^P(q^2)$:

$$\langle P(p_2) | \bar{q}\gamma^\mu(1 - \gamma^5)b | B(p_1) \rangle = p^\mu f_+^P(q^2) + q^\mu f_-^P(q^2), \quad (20)$$

where $p = p_1 + p_2$ and $q = p_1 - p_2$. For $B \rightarrow V\nu\bar{\nu}$ ($V = \rho, K^*$) decays, the hadronic matrix element can be written in terms of four form factors as

$$\begin{aligned} & \langle V(p_2, \varepsilon) | \bar{q}\gamma^\mu(1 - \gamma^5)b | B(p_1) \rangle \\ &= -\varepsilon_{\mu\nu\rho\sigma}\varepsilon^{*\nu}p_2^\rho q^\sigma \frac{2V^V(q^2)}{M_B + M_V} - i\left\{ \varepsilon_\mu^*(M_B + M_V)A_1^V(q^2) \right. \\ & \quad \left. - (\varepsilon^* \cdot q)(p_1 + p_2)_\mu \frac{A_2^V(q^2)}{M_B + M_V} - q_\mu(\varepsilon^* \cdot q) \frac{2M_V}{q^2}(A_3^V(q^2) - A_0^V(q^2)) \right\}, \quad (21) \end{aligned}$$

with $A_3^V(0) = A_0^V(0)$ and $A_3^V(q^2) = \frac{M_B + M_V}{2M_V}A_1^V(q^2) - \frac{M_B - M_V}{2M_V}A_2^V(q^2)$.

For the numerical analysis, we follow the theoretical estimates from LCSR model [32, 33]. The transition form factors are parameterized by three independent fit parameters and one

resonance mass. Form factors used in the numerical calculation are given by

$$f_+^\pi(q^2) = \frac{r_1}{1 - q^2/m_R^2} + \frac{r_2}{1 - q^2/m_{\text{fit}}^2}, \quad (22)$$

$$f_+^K(q^2) = \frac{r_1}{1 - q^2/m_R^2} + \frac{r_2}{(1 - q^2/m_R^2)^2}, \quad (23)$$

$$f_0^P(q^2) = \frac{r_2}{1 - q^2/m_{\text{fit}}^2}, \quad (24)$$

$$F^V(q^2) = \frac{r_1}{1 - q^2/m_R^2} + \frac{r_2}{1 - q^2/m_{\text{fit}}^2} \quad (\text{for } F = V \text{ or } A_0), \quad (25)$$

$$A_1^V(q^2) = \frac{r_2}{1 - q^2/m_{\text{fit}}^2}, \quad (26)$$

$$A_2^V(q^2) = \frac{r_1}{1 - q^2/m_{\text{fit}}^2} + \frac{r_2}{(1 - q^2/m_{\text{fit}}^2)^2}, \quad (27)$$

where values of parameters are shown in Table III and $f_-^P = (f_0^P - f_+^P) \frac{M_B^2 - M_P^2}{q^2}$.

REFERENCES

- [1] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Phys. Rev. Lett. **83**, 1914 (1999) [arXiv:hep-ph/9905312]; Nucl. Phys. B **591**, 313 (2000) [arXiv:hep-ph/0006124]; Nucl. Phys. B **606**, 245 (2001) [arXiv:hep-ph/0104110].
- [2] Y. Y. Keum, H. N. Li and A. I. Sanda, Phys. Rev. D **63**, 054008 (2001) [arXiv:hep-ph/0004173]; Phys. Lett. B **504**, 6 (2001) [arXiv:hep-ph/0004004].
- [3] C. W. Bauer, I. Z. Rothstein and I. W. Stewart, arXiv:hep-ph/0510241.
- [4] H. n. Li, S. Mishima and A. I. Sanda, Phys. Rev. D **72**, 114005 (2005) [arXiv:hep-ph/0508041].
- [5] S. Mishima and T. Yoshikawa, Phys. Rev. D **70**, 094024 (2004) [arXiv:hep-ph/0408090]; Y. L. Wu and Y. F. Zhou, *ibid.* **71**, 021701 (2005) [arXiv:hep-ph/0409221]; *ibid.*

- 72**, 034037 (2005) [arXiv:hep-ph/0503077]; A. J. Buras, R. Fleischer, S. Recksiegel and F. Schwab, Acta Phys. Polon. B **36** (2005) 2015 [arXiv:hep-ph/0410407]; arXiv:hep-ph/0512059; Y. Y. Charng and H. n. Li, Phys. Rev. D **71**, 014036 (2005) [arXiv:hep-ph/0410005]; X. G. He and B. H. J. McKellar, arXiv:hep-ph/0410098; S. Baek, P. Hamel, D. London, A. Datta and D. A. Suprun, Phys. Rev. D **71**, 057502 (2005) [arXiv:hep-ph/0412086]; C. S. Kim, S. Oh and C. Yu, *ibid.* **72**, 074005 (2005) [arXiv:hep-ph/0505060]; X. q. J. Li and Y. d. J. Yang, Phys. Rev. D **72**, 074007 (2005) [arXiv:hep-ph/0508079]; D. Chang, C. S. Chen, H. Hatanaka and C. S. Kim, arXiv:hep-ph/0510328.
- [6] R. Arnowitt, B. Dutta, B. Hu and S. Oh, Phys. Lett. B **633**, 748 (2006) [arXiv:hep-ph/0509233]. W. S. Hou, M. Nagashima and A. Soddu, Phys. Rev. Lett. **95**, 141601 (2005) [arXiv:hep-ph/0503072]; Phys. Rev. D **72**, 115007 (2005) [arXiv:hep-ph/0508237]; Y. D. Yang, R. Wang and G. R. Lu, *ibid.* **73**, 015003 (2006) [arXiv:hep-ph/0509273].
- [7] V. Barger, C. W. Chiang, P. Langacker and H. S. Lee, Phys. Lett. B **580**, 186 (2004) [arXiv:hep-ph/0310073]; *ibid.* **598**, 218 (2004) [arXiv:hep-ph/0406126]; C. H. Chen and H. Hatanaka, arXiv:hep-ph/0602140.
- [8] Y. Grossman, Z. Ligeti and E. Nardi, Nucl. Phys. B **465**, 369 (1996) [Erratum-*ibid.* B **480**, 753 (1996)] [arXiv:hep-ph/9510378].
- [9] G. Buchalla, G. Hiller and G. Isidori, Phys. Rev. D **63**, 014015 (2001) [arXiv:hep-ph/0006136].
- [10] T. M. Aliev and C. S. Kim, Phys. Rev. D **58**, 013003 (1998) [arXiv:hep-ph/9710428]; C. S. Kim, Y. G. Kim and T. Morozumi, *ibid.* **60**, 094007 (1999) [arXiv:hep-ph/9905528].
- [11] T. M. Aliev, A. Ozpineci and M. Savci, Phys. Lett. B **506**, 77 (2001) [arXiv:hep-ph/0101066].
- [12] C. Bobeth, A. J. Buras, F. Kruger and J. Urban, Nucl. Phys. B **630**, 87 (2002) [arXiv:hep-ph/0112305].

- [13] J. L. Hewett and T. G. Rizzo, Phys. Rept. **183**, 193 (1989).
- [14] A. Leike, Phys. Rept. **317**, 143 (1999) [arXiv:hep-ph/9805494].
- [15] J. Chay, K. Y. Lee and S. h. Nam, Phys. Rev. D **61**, 035002 (2000) [arXiv:hep-ph/9809298]. D. W. Jung, K. Y. Lee, H. S. Song and C. Yu, J. Korean Phys. Soc. **36**, 258 (2000) [arXiv:hep-ph/9905353]; K. Y. Lee, S. C. Park, H. S. Song and C. Yu, Phys. Rev. D **63**, 094010 (2001) [arXiv:hep-ph/0011173].
- [16] K. Leroux and D. London, Phys. Lett. B **526**, 97 (2002) [arXiv:hep-ph/0111246].
- [17] M. Cvetič and P. Langacker, Mod. Phys. Lett. A **11**, 1247 (1996) [arXiv:hep-ph/9602424].
- [18] G. Buchalla, G. Burdman, C. T. Hill and D. Kominis, Phys. Rev. D **53**, 5185 (1996) [arXiv:hep-ph/9510376]; G. Burdman, K. D. Lane and T. Rador, Phys. Lett. B **514**, 41 (2001) [arXiv:hep-ph/0012073].
- [19] G. Cleaver, M. Cvetič, J. R. Espinosa, L. L. Everett, P. Langacker and J. Wang, Phys. Rev. D **59**, 055005 (1999) [arXiv:hep-ph/9807479]; M. Cvetič, G. Shiu and A. M. Uranga, Phys. Rev. Lett. **87**, 201801 (2001) [arXiv:hep-th/0107143]; M. Cvetič, G. Shiu and A. M. Uranga, Nucl. Phys. B **615**, 3 (2001) [arXiv:hep-th/0107166]; M. Cvetič, P. Langacker and G. Shiu, Phys. Rev. D **66**, 066004 (2002) [arXiv:hep-ph/0205252].
- [20] M. Masip and A. Pomarol, Phys. Rev. D **60**, 096005 (1999) [arXiv:hep-ph/9902467].
- [21] N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, T. Gregoire and J. G. Wacker, JHEP **0208**, 021 (2002) [arXiv:hep-ph/0206020]; T. Han, H. E. Logan, B. McElrath and L. T. Wang, Phys. Rev. D **67**, 095004 (2003) [arXiv:hep-ph/0301040].
- [22] [ALEPH Collaboration], CERN-PPE-95-172 *Contributed to the 1995 Europhysics Conference on High Energy Physics - EPS-HEP, Brussels, Belgium, 27 Jul - 2 Aug 1995, and to the 17th International Symposium on Lepton-Photon Interactions, Beijing, China, 10 - 15 Aug 1995*
- [23] F. Abe *et al.* [CDF Collaboration], Phys. Rev. Lett. **77**, 438 (1996) [arXiv:hep-ex/9601008].

- [24] B. Abbott *et al.* [D0 Collaboration], FERMILAB-CONF-97-356-E *Presented at 18th International Symposium on Lepton and Photon Interactions (LP 97), Hamburg, Germany, 28 Jul - 1 Aug 1997, and Presented at International Europhysics Conference on High-Energy Physics (HEP 97), Jerusalem, Israel, 19-26 Aug 1997*
- [25] T. G. Rizzo, Phys. Rev. D **59**, 015020 (1999) [arXiv:hep-ph/9806397].
- [26] K. S. Babu, C. F. Kolda and J. March-Russell, Phys. Rev. D **54**, 4635 (1996) [arXiv:hep-ph/9603212]; *ibid.* **57**, 6788 (1998) [arXiv:hep-ph/9710441].
- [27] C. S. Kim, T. Morozumi, Sechul Oh and Chaehyun Yu, work in progress.
- [28] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996) [arXiv:hep-ph/9512380].
- [29] M. Misiak and J. Urban, Phys. Lett. B **451**, 161 (1999) [arXiv:hep-ph/9901278]; G. Buchalla and A. J. Buras, Nucl. Phys. B **548**, 309 (1999) [arXiv:hep-ph/9901288].
- [30] K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0507034.
- [31] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **94**, 101801 (2005) [arXiv:hep-ex/0411061].
- [32] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014015 (2005) [arXiv:hep-ph/0406232].
- [33] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014029 (2005) [arXiv:hep-ph/0412079].
- [34] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
- [35] P. Langacker and D. London, Phys. Rev. D **38**, 886 (1988).
- [36] T. Asaka and M. Shaposhnikov, Phys. Lett. B **620**, 17 (2005) [arXiv:hep-ph/0505013].